

STATISTICS IN TRANSITION *new series, March 2018*
Vol. 19, No. 1, pp. 7–23, DOI 10.21307/stattrans-2018-001

A BAYESIAN INFERENCE OF MULTIPLE STRUCTURAL BREAKS IN MEAN AND ERROR VARIANCE IN PANEL AR (1) MODEL

Varun Agiwal¹, Jitendra Kumar², Dahud Kehinde Shangodoyin³

ABSTRACT

This paper explores the effect of multiple structural breaks to estimate the parameters and test the unit root hypothesis in panel data time series model under Bayesian perspective. These breaks are present in both mean and error variance at the same time point. We obtain Bayes estimates for different loss function using conditional posterior distribution, which is not coming in a closed form, and this is approximately explained by Gibbs sampling. For hypothesis testing, posterior odds ratio is calculated and solved via Monte Carlo Integration. The proposed methodology is illustrated with numerical examples.

Key words: panel data model, autoregressive model, structural break, MCMC, posterior odds ratio.

1. Introduction

Statistical inference of panel data time series model received great attention in the last several decades in both econometrics and statistics literature. The main idea behind the use of panel data time series model is to overcome the difficulty of unobserved variation in cross-section data sets over individual units as well as variation, which may change the structure also. It was assumed that this change was taken by some observations at a fixed and common time point in each series referred to as break point. Thus, structural break concept in panel data set-up is important to handle the permanent effects in the series and impacts other simultaneous variables. For this, an extensive literature concentrates on testing, estimation and detection of the existence of single or multiple structural breaks from univariate to multivariate time series. Bai and Perron (1998, 2003) considered the problem of estimation and testing for break point in linear model and determine the number of breaks using double maximum tests. They have also further addressed various issues such as estimation and testing number of

¹ Department of Statistics, Central University of Rajasthan, Bandersindri, Ajmer, India.
E-mail: varunagiwal.stats@gmail.com.

² Department of Statistics, Central University of Rajasthan, Bandersindri, Ajmer, India.
E-mail: vjitendrav@gmail.com.

³ Department of Statistics, University of Botswana, Gaborone, Botswana.
E-mail: shangodoyink@gamil.com.

breaks, forming confidence interval related to multiple linear regression with multiple structural break. Altissimo and Corradi (2003) suggested an approach for detecting and estimation of the number of shifts in mean of a nonlinear process, which is having dependent and heterogeneous observations. They proposed a new estimator for long run variance, which was consistent in the presence of breaks and verified via a simulation exercise. Li (2004) applied quasi-Bayesian approach to detect the number and position of structural breaks in China's GDP and labour productivity data using predictive likelihood information criterion.

Apart from the above literature, which mainly dealt with the classical approach, a generalized form of estimation and testing the structural break by using Bayesian inference is less explored. Geweke and Jiang (2011) developed Bayesian approach to modelling in-sample structural breaks and forecasting out-of-sample breaks. Eo (2012) used Bayesian approach to estimate the number of breaks in autoregressive regressions with structural breaks in intercept, persistence, and residual variance. A model selection criterion was also considered to select the best model from U.S. GDP deflator data. Aue and Horvath (2013) discussed several approaches for estimating the parameter and locating multiple break points. They considered CUSUM procedure as well as likelihood statistic to adjust the serial dependence in presence of structural break. Recently, Melighotsiduo *et al.* (2017) suggested a Bayesian approach for autoregressive model allowing multiple structural changes in both mean and error variance of economic series occurring at unknown times, and Bayesian unit root testing is also proposed.

In current scenario, a growing literature on estimation and testing of multiple structural breaks in generalized univariate model such as panel data as well as multivariate time series model. A partial list of contributions in multiple structural breaks include Sugita (2006), Liu *et al.* (2011), Jin *et al.* (2013), Preuss *et al.* (2015) and Eo and Morley (2015) to analysis the procedure for detection and estimation of change point in vector error correction model, panel data model. In recent time, detection and estimation of multiple change points in panel data with interactive fixed effect and dynamic structure is introduced. Li *et al.* (2016) through penalized principal component (PPC) estimation procedure with an adaptive group fused LASSO.

An overview of the above description, this paper provides a general methodology to estimate and inference for panel data model under the presence of multiple change points in mean and error variance parameters. Our approach provides a flexible way to the interpretation of the result in real situation because in most economic and time series data are varying by trend and variance component. If one considers a break in mean also, then the impact of the series changes due to both type of break versus no break point. Thus, a Bayesian approach is introduced to capture the impact of break points in the panel data model. For Bayes estimation, we apply both symmetric and asymmetric loss function to posterior density in order to get better estimators and compare them with ordinary least square estimator. In addition, we also examine the model selection criterion to find the appropriate model, which may or may not contain multiple break points in a real data set.

2. Model Specification

Let $\{y_{it}, t=1,2,\dots,T; i=1,2,\dots,n\}$ be a panel data time series model having B multiple break points in mean and error variance where breaks occur in both parameters in the same location. In that case our panel data model can be expressed as

$$y_{it} = \begin{cases} \rho y_{i,t-1} + (1-\rho)\mu_{i1} + \sigma_1 \varepsilon_{it}; & T_0 < t \leq T_1 \\ \vdots & \vdots \\ \rho y_{i,t-1} + (1-\rho)\mu_{ij} + \sigma_j \varepsilon_{it}; & T_{j-1} < t \leq T_j \\ \vdots & \vdots \\ \rho y_{i,t-1} + (1-\rho)\mu_{i,B+1} + \sigma_{B+1} \varepsilon_{it}; & T_B < t \leq T_{B+1} = T \end{cases} \quad (1)$$

for $j = 1,2,\dots, B$ and where n denotes number of cross sectional units, ρ is the autoregressive coefficient, μ_j is a $(n \times 1)$ vector of mean coefficients at j^{th} division and ε_{it} are assumed to be independent and normally distributed with zero mean and division specific variance σ_j^2 . This is a partial structural change model since the parameter ρ is not subject to shifts and is estimated using the entire sample space. The model in (1) can also be casted in the form of matrix notation with \cdot^* Kronecker delta product indicating element by element array multiplication, Z as the $nT \times (B+1)$ matrix whose j^{th} column is equal to one if $T_{j-1} < t \leq T_j$ and zero otherwise, and consider mean and residual variance parameters as a vector form.

$$y = \rho y_{-1} + (1-\rho)L\mu + \varepsilon ; \quad \varepsilon \sim N(0, I_{nT} \cdot^* S)$$

where

$$y = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}_{nT \times 1}; y_i = \begin{pmatrix} y_{i1} \\ y_{i2} \\ \vdots \\ y_{iT} \end{pmatrix}_{T \times 1}; \mu = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_{B+1} \end{pmatrix}_{n(B+1) \times 1}; \mu_j = \begin{pmatrix} \mu_{1j} \\ \mu_{2j} \\ \vdots \\ \mu_{nj} \end{pmatrix}_{n \times 1}; \sigma = \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \vdots \\ \sigma_{B+1} \end{pmatrix}_{(B+1) \times 1}$$

$$L = \begin{pmatrix} I_n \otimes l_{T_1} & 0 & \dots & 0 \\ 0 & I_n \otimes l_{T_2-T_1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & I_n \otimes l_{T-T_B} \end{pmatrix}; Z = \begin{pmatrix} l_{T_1} & 0 & \dots & 0 \\ 0 & l_{T_2-T_1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & l_{T-T_B} \end{pmatrix}; S = Z\sigma$$

Our study attempts to estimate the parameters in structural break model under Bayesian framework and test the unit root hypothesis by using posterior odds ratio. Under unit root case, model (1) reduces to a pure structural change model where all the model's coefficients are subject to change

$$\Delta y_{it} = \begin{cases} \sigma_1 \varepsilon_{it}; & T_0 < t \leq T_1 \\ \vdots & \vdots \\ \sigma_j \varepsilon_{it}; & T_{j-1} < t \leq T_j \\ \vdots & \vdots \\ \sigma_{B+1} \varepsilon_{it}; & T_B < t \leq T_{B+1} \end{cases} \quad (2)$$

As mentioned, if we follow the usual approach defined in the literature to test for stationarity model reduces by (2) under the null hypothesis $H_0: \rho = 1$ is difference stationary with multiple breaks in error variance against the alternative hypothesis $H_1: \rho \in S$, series is stationary with multiple breaks in mean as well as error variance.

3. Bayesian Inference

In this section, we discuss issues related to the estimation and inference about the parameters and testing of unit root hypothesis. In order to perform Bayesian inference we need the likelihood function and specify prior distribution for the model parameters. Posterior probability is obtained by using sample information contained in the likelihood function combined with the joint prior distribution. The likelihood function for this model is

$$L(\rho, \mu, \sigma | y) = (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \left(\sigma_j^{-n(T_j - T_{j-1})} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1} - (1-\rho)\mu_{ij})^2 \right\} \right] \quad (3)$$

For panel data model generally normal prior distribution is considered for $\mu_{ij} \sim N(\gamma_{ij}, \sigma_j^2)$, for error variance (σ_j^2) assume conjugate inverted gamma prior $IG(c_j, d_j)$ and uniform prior is taken for autoregressive coefficient (ρ), see [Schotman and Van Dijk (1991) and Phillips (1991)]. The joint prior distribution is given as

$$\pi(\rho, \mu, \sigma^2) = \frac{(2\pi)^{-\frac{n(B+1)}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} (\sigma_j^2)^{-c_j - \frac{n}{2}} \right) \exp \left[-\sum_{j=1}^{B+1} \frac{1}{\sigma_j^2} \left\{ d_j + \frac{1}{2} \sum_{i=1}^n (\mu_{ij} - \gamma_{ij})^2 \right\} \right] \quad (4)$$

3.1. Bayesian Estimation via Gibbs Sampling

Given the likelihood function and prior density defined by eqⁿ (3) and eqⁿ (4), the posterior distribution is given by

$$\pi(\rho, \mu, \sigma | y) \propto \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1} - (1-\rho)\mu_{ij})^2 + \sum_{i=1}^n (\mu_{ij} - \gamma_{ij})^2 + 2d_j \right) \right\} \right] \tag{5}$$

The posterior distribution in (5) is very complicated and hence no closed form inference appears to be possible. For Bayesian estimation, we proceed via Gibbs sampler, a MCMC method, proposed by Geman and Geman (1984). The Gibbs sampler procedure, which we used, is described by Wang and Zivot (2000) in a time series regression model with multiple structural breaks. By means of this procedure, it gives a chain of estimated parameters values, which is frequently obtained by conditional probability distribution. Here, our aim is to generate a sequence of random variables from the conditional probability distribution using the current value of the parameters. For this we have derived the form of conditional posterior distributions given below:

$$\pi(\mu_{ij} | \rho, \sigma_j^2, \underline{y}) \sim N \left(\frac{(1-\rho) \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1}) + \gamma_{ij}}{(1-\rho)^2 (T_j - T_{j-1}) + 1}, \frac{\sigma_j^2}{(1-\rho)^2 (T_j - T_{j-1}) + 1} \right) \tag{6}$$

$$\pi(\sigma_j^2 | \rho, \mu_{ij}, \underline{y}) \sim IG \left(\frac{n(T_j - T_{j-1}) + n}{2} + c_j, d_j + \frac{1}{2} \sum_{i=1}^n \left(\sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1} - (1-\rho)\mu_{ij})^2 + (\mu_{ij} - \gamma_{ij})^2 \right) \right) \tag{7}$$

$$\pi(\rho | \mu_{ij}, \sigma_j^2, \underline{y}) \sim TN \left(\frac{\sum_{j=1}^{B+1} \sum_{t=T_{j-1}}^{T_j} (y_{it} - \mu_{ij})(y_{i,t-1} - \mu_{ij})}{\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it-1} - \mu_{ij})^2}, \frac{\sum_{j=1}^{B+1} \sigma_j^2}{\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it-1} - \mu_{ij})^2}, l, 1 \right) \tag{8}$$

Using the generated samples from the above posteriors, Bayes estimates of the parameter are evaluated by different loss functions under Gibbs sampling algorithm. A loss function is a decision rule to select the best estimator and represent each of the possible estimates. Here, we consider squared error (symmetric) loss function as well as entropy (asymmetric) loss function for getting better understanding of the Bayesian estimation. Under squared error and entropy loss function, Bayes estimator are $E(\theta | \underline{x})$ and $[E(\theta^{-1} | \underline{x})]^{-1}$.

3.2. Testing Unit Root Hypothesis via Posterior Odds Ratio

In a hypothesis testing problem, one is generally interested in testing the stationary condition of a model. Here, null hypothesis is used as a unit root hypothesis against the alternative of a stationary model. In Bayesian framework, testing is often convenient to summarize the information in terms of posterior odds ratio. The posterior odds ratio is the ratio of posterior probability under null versus alternative hypothesis with the product of prior odds, notation given as:

$$\beta_{01} = \frac{p_0}{1-p_0} \frac{P(y|H_0)}{P(y|H_1)} \tag{9}$$

Theorem: To test the null hypothesis that y_{it} is a non-stationary I(1) process, i.e. $\rho=1$ in equation (2), against the alternative hypothesis that y_{it} is a stationary I(0) process, i.e. $\rho \in S$ in equation (1). The posterior odds ratio can be constructed according to equation (9)

$$\beta_{01} = \frac{p_0}{1-p_0} \frac{\prod_{j=1}^{B+1} \left[d_j + \frac{1}{2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - y_{i,t-1})^2 \right]^{\frac{n(T_j - T_{j-1}) + 2c_j}{2}}}{\frac{1}{1-l} \int \prod_{l=1}^{B+1} [A_j(\rho)]^{\frac{n}{2}} [C_j(\rho)]^{\frac{n(T_j - T_{j-1}) + 2c_j}{2}} d\rho} \tag{10}$$

where

$$A_j(\rho) = (1-\rho)^2 (T_j - T_{j-1}) + 1$$

$$B_{ij}(\rho) = (1-\rho) \sum_{t=T_{j-1}}^{T_j} (y_{it} - y_{i,t-1}) + \gamma_{ij}$$

$$C_j(\rho) = d_j + \frac{1}{2} \sum_{i=1}^n \left(\sum_{t=T_{j-1}}^{T_j} (y_{it} - y_{i,t-1})^2 + \gamma_{ij}^2 - \frac{[B_{ij}(\rho)]^2}{A_j(\rho)} \right)$$

Proof: The proof of the theorem is given in the appendix.

In the equation (10), closed form expression of posterior odds ratio is not obtained. Therefore, we use an alternative technique as Monte Carlo integration for approximately solving the integrals and get the value of posterior odds ratio.

4. Simulation Study

In this section, we conduct a set of simulated experiments to evaluate the performance of our model and compare different estimators based on Monte Carlo simulation. To estimate the model parameters, assume that the number of breaks and the location of break points are known so that the remaining objectives in equation (1) are estimated via an iterative procedure. In simulation experiment we have generated artificial time series from our model with varying

numbers of structural breaks at the same time points in mean and error variance parameters. We are starting with the initial observation $y_{0i} = (10, 15, 20)$ to generate panel data time series from the suggested model having three panel ($n=3$) and each panel contains T observations. For better interpretation, we took different size of time series $T = (50, 75, 100)$ and also varying autoregressive coefficients $\rho = (0.9, 0.95, 0.99)$. The number of possible structural break (B) has been 3. Thus, the disturbances ϵ_{it} are generated as i.i.d. for all i and j with four different variance, namely $(\sigma_1^2, \sigma_2^2, \sigma_3^2, \sigma_4^2) = (0.1, 0.2, 0.3, 0.4)$. For inverse gamma prior distribution with hyper parameters is to known. For numerical purpose we have taken as $c_j = d_j = 0.01$ for all break points. In the case of normal prior, hyper prior mean is equal to mean of the generated series at every break point interval (T_{j-1}, T_j) with parallel variance given in disturbances term. The true value of mean term for each panel having four partitions is written as $(\mu_{11}, \mu_{12}, \mu_{13}) = (14, 16, 18)$; $(\mu_{21}, \mu_{22}, \mu_{23}) = (20, 22, 24)$; $(\mu_{31}, \mu_{32}, \mu_{33}) = (26, 28, 30)$; $(\mu_{41}, \mu_{42}, \mu_{43}) = (32, 34, 36)$. All results are based on 5000 replications. From the generated sample, we obtained Bayes estimate of parameters and compared the performance with ordinary least square (OLS) estimate. We report the estimated value and its mean square error in Table-4.1 to 4.3.

Table-4.1. Estimators with varying time series at $\rho = 0.9$

	T=50			T=75			T=100		
	OLS	SELF	ELF	OLS	SELF	ELF	OLS	SELF	ELF
ρ	0.8607	0.9037	0.9036	0.8724	0.9049	0.9048	0.8763	0.9043	0.9043
	0.0025	0.0000	0.0000	0.0012	0.0001	0.0001	0.0009	0.0001	0.0001
μ_{11}	13.7381	13.9853	13.9785	13.8519	13.9959	13.9893	13.7567	13.9815	13.9749
	0.5840	0.0104	0.0107	0.5529	0.0098	0.0100	0.5706	0.0091	0.0094
μ_{12}	16.1184	15.9989	15.9929	16.0870	15.9938	15.9880	16.1338	16.0026	15.9969
	0.7121	0.0124	0.0125	0.5515	0.0106	0.0107	0.6236	0.0110	0.0111
μ_{13}	18.2558	17.9770	17.9717	18.2331	17.9833	17.9782	18.2917	17.9964	17.9912
	0.7394	0.0104	0.0106	0.7099	0.0108	0.0110	0.7042	0.0100	0.0101
μ_{21}	18.9938	20.0110	20.0014	19.4758	20.0186	20.0098	19.6677	20.0093	20.0015
	2.6156	0.0100	0.0099	1.2051	0.0235	0.0234	0.6942	0.0386	0.0385
μ_{22}	20.8791	21.9925	21.9838	21.4667	22.0115	22.0035	21.7912	22.0295	22.0224
	2.6572	0.0151	0.0154	1.5227	0.0293	0.0292	0.6849	0.0359	0.0356
μ_{23}	23.1734	24.0115	24.0035	23.5620	24.0168	24.0094	23.7996	24.0274	24.0209
	2.2536	0.0116	0.0114	1.2840	0.0278	0.0275	0.5568	0.0325	0.0322
μ_{31}	24.9505	26.0408	26.0295	25.5998	26.0509	26.0410	25.7780	26.0333	26.0244
	4.4370	0.0257	0.0248	1.7431	0.0381	0.0373	0.7004	0.0381	0.0376
μ_{32}	26.7468	28.0131	28.0026	27.6054	28.0537	28.0445	27.7914	28.0448	28.0364
	4.8589	0.0230	0.0228	1.7379	0.0385	0.0377	0.9369	0.0557	0.0550
μ_{33}	28.6660	30.0059	29.9961	29.5123	30.0316	30.0230	29.6804	30.0104	30.0027
	5.1925	0.0218	0.0218	1.7849	0.0359	0.0355	0.8159	0.0444	0.0443
μ_{41}	30.9830	32.0348	32.0234	31.1368	31.9948	31.9832	31.4541	32.0210	32.0093
	3.6570	0.0363	0.0358	3.9274	0.0448	0.0450	2.8567	0.0384	0.0382
μ_{42}	32.9950	34.0339	34.0232	33.5021	34.0500	34.0391	33.6176	34.0352	34.0243
	4.4146	0.0376	0.0369	2.5846	0.0370	0.0361	2.8885	0.0380	0.0374
μ_{43}	34.8637	36.0233	36.0132	35.1567	36.0032	35.9929	35.4250	36.0242	36.0139
	3.7348	0.0342	0.0340	3.4651	0.0386	0.0388	2.5830	0.0353	0.0349
σ_1^2	0.1071	0.1085	0.0988	0.1042	0.1053	0.1014	0.1038	0.1046	0.1005
	0.0007	0.0007	0.0000	0.0004	0.0005	0.0000	0.0005	0.0005	0.0000
σ_2^2	0.2104	0.2094	0.1961	0.2065	0.2069	0.2052	0.2031	0.2037	0.2030
	0.0040	0.0034	0.0000	0.0017	0.0015	0.0001	0.0006	0.0006	0.0000
σ_3^2	0.3121	0.3189	0.3028	0.3002	0.3031	0.2920	0.3054	0.3060	0.2984
	0.0075	0.0071	0.0003	0.0034	0.0034	0.0001	0.0018	0.0018	0.0000
σ_4^2	0.4058	0.4116	0.4027	0.4140	0.4193	0.3987	0.4252	0.4241	0.4221
	0.0055	0.0055	0.0004	0.0078	0.0078	0.0002	0.0100	0.0094	0.0011

Table-4.2. Estimators with varying time series at $\rho = 0.95$

	T=50			T=75			T=100		
	OLS	SELF	ELF	OLS	SELF	ELF	OLS	SELF	ELF
ρ	0.9277	0.9511	0.9511	0.9239	0.9511	0.9510	0.9231	0.9515	0.9515
	0.0010	0.0000	0.0000	0.0012	0.0000	0.0000	0.0011	0.0000	0.0000
μ_{11}	14.2813	14.0055	13.9979	14.3639	14.0031	13.9955	14.2784	14.0008	13.9934
	2.8644	0.0026	0.0026	2.8140	0.0026	0.0027	2.4136	0.0021	0.0021
μ_{12}	17.0431	16.0055	15.9989	16.8423	15.9981	15.9916	17.0321	16.0007	15.9942
	4.9016	0.0027	0.0027	4.3718	0.0029	0.0029	3.7536	0.0022	0.0022
μ_{13}	19.2725	17.9970	17.9910	19.2403	17.9869	17.9811	19.5766	17.9947	17.9889
	5.8300	0.0026	0.0027	5.7341	0.0021	0.0022	6.0293	0.0021	0.0022
μ_{21}	19.1892	20.0027	19.9902	19.1421	20.0079	19.9981	19.0485	20.0086	19.9988
	18.5998	0.0029	0.0030	6.4266	0.0088	0.0087	3.9955	0.0082	0.0082
μ_{22}	21.1867	21.9990	21.9876	21.3103	22.0017	21.9928	21.2755	21.9997	21.9907
	10.5883	0.0025	0.0027	4.6515	0.0079	0.0080	3.2040	0.0079	0.0080
μ_{23}	23.8031	24.0070	23.9966	23.6473	24.0044	23.9962	23.7322	24.0062	23.9980
	11.7370	0.0029	0.0028	3.8351	0.0074	0.0074	3.0148	0.0089	0.0088
μ_{31}	23.5217	25.9969	25.9830	23.9895	25.9922	25.9803	24.5652	26.0025	25.9912
	30.2210	0.0041	0.0043	9.4940	0.0073	0.0076	6.4169	0.0145	0.0146
μ_{32}	26.4880	28.0077	27.9947	26.7673	28.0165	28.0056	26.7426	28.0081	27.9977
	22.0991	0.0037	0.0037	9.3507	0.0119	0.0117	5.5741	0.0151	0.0151
μ_{33}	28.4393	29.9977	29.9856	28.7290	30.0130	30.0027	28.7848	30.0002	29.9904
	25.9163	0.0038	0.0039	7.4535	0.0120	0.0119	5.2416	0.0146	0.0148
μ_{41}	29.9873	31.9863	31.9745	30.8228	32.0145	32.0025	31.0121	32.0101	31.9987
	10.5213	0.0195	0.0201	8.4309	0.0336	0.0335	4.5547	0.0305	0.0305
μ_{42}	32.9922	34.0512	34.0402	32.9787	34.0263	34.0151	33.0487	34.0116	34.0008
	8.6453	0.0235	0.0224	5.3532	0.0246	0.0242	4.1663	0.0297	0.0296
μ_{43}	34.7339	36.0139	36.0035	35.2679	36.0439	36.0333	35.0919	36.0133	36.0032
	8.8365	0.0193	0.0192	5.1718	0.0286	0.0278	4.1124	0.0292	0.0291
σ_1^2	0.1087	0.1058	0.1019	0.1077	0.1048	0.1017	0.1065	0.1047	0.0999
	0.0011	0.0010	0.0000	0.0007	0.0007	0.0000	0.0008	0.0008	0.0000
σ_2^2	0.2566	0.2515	0.2244	0.1889	0.2040	0.1978	0.2057	0.2030	0.2000
	0.0126	0.0100	0.0019	0.0016	0.0017	0.0002	0.0015	0.0014	0.0000
σ_3^2	0.3739	0.3679	0.3372	0.3182	0.3137	0.2980	0.3094	0.3062	0.3036
	0.0265	0.0211	0.0016	0.0049	0.0048	0.0003	0.0026	0.0026	0.0000
σ_4^2	0.3961	0.4001	0.3946	0.4132	0.4091	0.4032	0.3940	0.4016	0.3984
	0.0043	0.0043	0.0002	0.0035	0.0034	0.0001	0.0024	0.0024	0.0000

Table-4.3. Estimators with varying time series at $\rho = 0.99$

	T=50			T=75			T=100		
	OLS	SELF	ELF	OLS	SELF	ELF	OLS	SELF	ELF
ρ	0.9887	0.9900	0.9900	0.9896	0.9900	0.9900	0.9896	0.9900	0.9900
	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_{11}	14.0541	13.9998	13.9995	13.8541	13.9996	13.9993	13.9496	13.9991	13.9988
	2.5280	0.0000	0.0000	2.2032	0.0000	0.0000	1.4473	0.0000	0.0000
μ_{12}	15.7119	15.9994	15.9992	16.1693	16.0004	16.0002	16.1484	16.0006	16.0004
	2.7599	0.0000	0.0000	1.7276	0.0000	0.0000	1.9185	0.0000	0.0000
μ_{13}	18.0323	18.0016	18.0014	17.9392	18.0001	17.9998	18.0924	18.0009	18.0006
	2.8263	0.0000	0.0000	2.4351	0.0001	0.0001	2.0293	0.0000	0.0000
μ_{21}	20.0894	20.0014	20.0011	20.1550	20.0008	20.0005	19.7357	19.9988	19.9985
	15.3991	0.0001	0.0001	5.6088	0.0001	0.0001	4.1294	0.0001	0.0001

Table-4.3. Estimators with varying time series at $\rho = 0.99$ (cont.)

	T=50			T=75			T=100		
	OLS	SELF	ELF	OLS	SELF	ELF	OLS	SELF	ELF
μ_{22}	22.1373	21.9999	21.9996	21.8658	22.0008	22.0005	22.0746	22.0001	21.9998
	17.3947	0.0001	0.0001	6.3083	0.0001	0.0001	3.4212	0.0001	0.0001
μ_{23}	24.1136	24.0001	23.9998	23.8898	23.9995	23.9992	24.1808	24.0002	23.9999
	20.6617	0.0001	0.0001	3.6968	0.0001	0.0001	4.7187	0.0001	0.0001
μ_{31}	25.4894	26.0006	26.0002	25.9236	26.0004	26.0001	25.8274	25.9991	25.9988
	50.4575	0.0001	0.0001	9.9667	0.0001	0.0001	9.8900	0.0001	0.0001
μ_{32}	27.7662	27.9993	27.9991	27.8165	28.0000	27.9997	27.5075	28.0013	28.0010
	37.2489	0.0001	0.0001	15.0037	0.0001	0.0001	11.5979	0.0001	0.0001
μ_{33}	29.7095	30.0006	30.0003	29.7752	30.0009	30.0006	30.0822	29.9988	29.9985
	36.9404	0.0001	0.0001	12.4540	0.0001	0.0001	7.6263	0.0001	0.0001
μ_{41}	31.7953	32.0001	31.9998	32.1870	32.0006	32.0004	31.7316	31.9987	31.9985
	72.6862	0.0001	0.0001	7.7369	0.0001	0.0001	2.9958	0.0001	0.0001
μ_{42}	33.7731	34.0002	33.9999	33.9305	34.0000	33.9998	33.8511	34.0027	34.0024
	72.3302	0.0001	0.0001	6.8616	0.0001	0.0001	2.8177	0.0001	0.0001
μ_{43}	35.9221	35.9988	35.9985	35.9559	35.9998	35.9995	35.8189	36.0004	36.0002
	75.6086	0.0001	0.0001	6.9744	0.0001	0.0001	2.9922	0.0001	0.0001
σ_1^2	0.1191	0.1088	0.1006	0.1221	0.1121	0.0993	0.1153	0.1122	0.1017
	0.0024	0.0020	0.0000	0.0034	0.0017	0.0003	0.0024	0.0016	0.0001
σ_2^2	0.2396	0.2082	0.1962	0.2306	0.2288	0.1845	0.2405	0.2236	0.2025
	0.0088	0.0077	0.0002	0.0097	0.0093	0.0003	0.0122	0.0071	0.0001
σ_3^2	0.3452	0.3208	0.2999	0.3418	0.3215	0.2642	0.3502	0.3313	0.2817
	0.0172	0.0123	0.0002	0.0227	0.0139	0.0018	0.0253	0.0207	0.0009
σ_4^2	0.3953	0.4094	0.3891	0.3920	0.4080	0.4013	0.4067	0.4005	0.4011
	0.0036	0.0032	0.0004	0.0017	0.0018	0.0000	0.0012	0.0012	0.0001

Table-4.1-4.3 shows the behaviour of ordinary least square (OLS) and Bayes estimators of parameter with varying values of time series at different autoregressive coefficient. It can be easily seen that MSEs of all estimators decrease as the sample size of series increases. The difference between the estimated value and the true value in OLS is large, which explains that average bias is maximum as compared to Bayes estimator. The Bayes estimator under both loss functions performs better because of additional information given about the parameter. A better estimated value for autoregressive coefficient, mean term and error variance is obtained by ELF as compared to SELF due to less MSE for low value of ρ . For a high value of ρ , Bayes estimator obtained under different loss function is equally applicable to estimate the parameters since both the estimators show more or less same magnitudes for their MSE. An increase in the number of breaks points, MSE also increases because of the length of the segment is small and take less observation to estimate the parameters.

After estimation of structural break parameters, testing of the unit root is considered. We can calculate posterior odds ratio values with different values of ρ and varying size of the series, which are reported in Table-4.4. The table shows

that as the value of ρ increases, POR reduces to zero and this rejects the null hypothesis, i.e. unit root hypothesis. Thus, the model which contains multiple breaks in mean and variance have a stationary model for this simulated series.

Table-4.4. Posterior odds ratio with varying ρ and T

T	$\rho=0.90$	$\rho=0.92$	$\rho=0.94$	$\rho=0.96$	$\rho=0.98$
50	0.0750	0.0416	0.0245	0.0098	0.0037
60	0.0645	0.0412	0.0291	0.0118	0.0044
70	0.0723	0.0447	0.0316	0.0149	0.0060
80	0.0647	0.0441	0.0338	0.0159	0.0067

5. Real Data Analysis

To provide a practical application of our model and verify the result obtained by simulation study, we apply our proposed work to a real data set. We use agricultural production and productivity of various crops of food grains data set consisting of 60 years' time series of Rice (R), Wheat (W) and Coarse Cereals (CC) variables for the annually book of "*Handbook of Statistics on Indian Economy*" from 1954-55 to 2014-15. The source of food grains data set is taken through Ministry of Agriculture & Farmers Welfare, Government of India by Reserve Bank of India and the book was published by Data Management and Dissemination Division (DMDD), Department of Statistics and Information Management (DSIM), Reserve Bank of India (RBI). This book provides statistical data on a wide range of economic and financial indicators related to national income variable, output, prices, money, banking, financial markets, etc. To determine the number of break points and their positions in food grains data set, we can use "*strucchange*" package developed by Zeileis *et al.* (2002) in R-language. The command "*breakpoints*" is considered to suggest the number of break points and identify their location in respective individual series. The results are reported in Table-5.1, which is given below:

Table-5.1: Structural break point present in different cereals

Break Point	Rice	Wheat	Coarse Cereals
T ₁	12	12	12
T ₂	22	22	33
T ₃	33	30	51
T ₄	42	39	
T ₅	51	51	

From the above table one can observe that each series could not contain equal number of break points and their positions also differ from one series to another because of finding the break points individually. After applying the procedure we observe that each series includes two similar break points 12 and 51, which is near and far from the series. The remaining break points (22, 30, 33, 39, 42) mainly occur in between these points. For analysis, we make different

combinations of break points containing various numbers of breaks. There are seven break points established; 127 combinations included single as well as multiple breaks. To identify a suitable model for this data set by using log-likelihood function, Akaike information criterion (AIC) and Bayes information criterion (BIC) define how many break points and their location is present in the model. For examination, consider only starting eight combinations, which have minimum AIC and BIC values, as shown in the Table-5.2.

Table-5.2. Break point detection

Break point	POR	ρ	AIC	BIC	Log L
(22,42,51)	0.00878	0.9803	1433.0263	1474.3162	-703.5131
(22,30,51)	0.01617	0.9778	1454.7726	1496.0625	-714.3863
(22,30,42,51)	0.00362	0.9732	1460.8775	1511.6959	-714.4387
(22,42)	0.06997	0.9808	1471.8446	1513.6061	-725.9223
(12,30,51)	0.01872	0.9749	1477.9163	1519.2062	-725.9581
(12,30,42,51)	0.00745	0.9693	1487.6784	1538.4968	-727.8392
22	0.51915	0.9881	1524.1465	1546.3796	-755.0733
(22,30,42)	0.01195	0.9738	1524.5874	1565.8773	-749.2937

By using information criterion, Table-5.2 gives appropriate conclusion about the number of break points and their positions to obtain a suitable model for this data set. The table shows that data follow a PAR(1) model having three break points (22, 42, 51) because of minimum AIC and BIC. Considering higher number of break points, i.e. 5, 6 or 7 in the series, no combination occurs in the last eight observations because the increase in the break point is inconvenient to partition the series in small segments. The maximum number of break points is 4 in this table, which can be considered after 3 break points. When the break point position occurs at only single point, i.e. 22, then AIC and BIC have large value as compared to two break points (22, 42). If we think about these two break points, AIC and BIC values may or may not be larger than three or four break points at different locations. If positions occur mostly at 22 and 51, statistic values is minimum in our combination so that mean 7C_i , $i = 1$ to 7. Minimum difference between break points may increase the statistic values, which directly reject these points of the model. An increase in the number of break points, posterior odds ratio value tends to zero, which concludes our model, which contains multiple breaks in mean and variance, is a better model compared to no-break model, i.e. PAR(1) model. The table also shows that data series is a stationary series for any combination of break point considered in the model. As a break point increases, the value of POR tends to zero, which concludes the model is stationary and no unit root is present in the model to make this as a difference stationary.

Once we acquire the number of break points and their positions in the proposed model, use this to estimate the parameters of the model for the data set using Gibbs procedure. The results are summarized in the Table-5.3. The table gives an appropriate value of the estimate parameter for the food grains data set in different types of estimation technique.

Table-5.3. Estimates Value Using Real Data Set

Parameter	OLS	SELF	ELF
ρ	0.7786	0.7788	0.7788
μ_{11}	42.6226	41.6685	41.6810
μ_{12}	21.3355	20.8544	20.8631
μ_{13}	26.8882	26.2820	26.2926
μ_{21}	72.9170	71.2050	71.2258
μ_{22}	57.7628	56.4043	56.4220
μ_{23}	31.0778	30.3408	30.3538
μ_{31}	94.2339	91.1963	91.2235
μ_{32}	77.5137	75.0114	75.0358
μ_{33}	34.8718	33.7356	33.7521
μ_{41}	109.4301	105.7798	105.8104
μ_{42}	95.1015	91.9243	91.9550
μ_{43}	46.2301	44.6730	44.6951
σ_1^2	12.9941	12.9820	12.9262
σ_2^2	25.5067	25.4830	25.3736
σ_3^2	51.1556	51.1081	50.8886
σ_4^2	25.9380	25.9139	25.8027

6. Conclusions

This paper deals with multiple structural breaks, which are present in mean and error variance in panel AR (1) model. Bayesian framework is used for estimating and testing the unit root hypothesis. Bayesian estimator gives better estimated value of the parameter as compared to OLS estimator in simulation as well as in real data. Testing of the hypothesis gives appropriate conclusion about the simulated series, which is stationary, and this is also verified by the real data set at each combination of break points. Break point identification is also done in real data set by using information criterion. This model may be extended to panel AR (p) model with similar types of breaks as well as to VAR model.

Acknowledgement

The second author gratefully acknowledges the financial assistance from UGC, India under MRP Scheme (Grant No.42-43/2013).

REFERENCES

- ALTISSIMO, F., CORRADI, V., (2003). Strong rules for detecting the number of breaks in a time series. *Journal of Econometrics*, 117 (2), pp. 207–244.
- AUE, A., HORVÁTH, L., (2013). Structural breaks in time series. *Journal of Time Series Analysis*, 34 (1), pp. 1–16.
- BAI, J., PERRON, P., (1998). Estimating and testing linear models with multiple structural changes. *Econometrica*, 66 (1), pp. 47–78.
- BAI, J., PERRON, P., (2003). Computation and analysis of multiple structural change models. *Journal of Applied Econometrics*, 18 (1), pp. 1–22.
- EO, Y., MORLEY, J., (2015). Likelihood-based confidence sets for the timing of structural breaks. *Quantitative Economics*, 6, pp. 463–497.
- EO, Y., (2012). Bayesian Inference about the Types of Structural Breaks When There are Many Breaks. Available at SSRN: <http://dx.doi.org/10.2139/ssrn.2011825>
- GEMAN, S., GEMAN, D., (1984). Stochastic relaxation, Gibbs distribution and the Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 6, pp. 721–741.
- GEWEKE, J., JIANG, Y., (2011). Inference and prediction in a multiple-structural-break model. *Journal of Econometrics*, 163 (2), pp. 172–185.
- JIN, B., SHI, X., WU, Y., (2013). A novel and fast methodology for simultaneous multiple structural break estimation and variable selection for nonstationary time series models. *Statistics and Computing*, 23 (2), pp. 1–11.
- LI, D., QIAN, J., SU, L., (2016). Panel data models with interactive fixed effects and multiple structural breaks. *Journal of the American Statistical Association*, 111 (516), pp. 1804–1819.
- LI, X., (2004). A Quasi-Bayesian Analysis of Structural Breaks in China's Output and Productivity Series. *International Journal of Business and Economics*, 3 (1), pp. 57–65.
- LIU, D., LI, R., WANG, Z., (2011). Testing for structural breaks in panel varying coefficient models: with an application to OECD health expenditure. *Empirical Economics*, 40 (1), pp. 95–118.
- MELIGKOTSIDOU, L., TZAVALIS, E., VRONTOS, I. D., (2017). On Bayesian Analysis and Unit Root Testing for Autoregressive Models in the Presence of Multiple Structural Breaks. *Econometrics and Statistics*, 4, pp.70–90.
- PHILLIPS, P.C., (1991) To criticize the critics: An objective Bayesian analysis of stochastic trends. *Journal of Applied Econometrics*, 6, pp.333–364.
- PREUSS, P., PUCHSTEIN, R., DETTE, H., (2015). Detection of multiple structural breaks in multivariate time series. *Journal of the American Statistical Association*, 110 (510), pp. 654–668.

- SCHOTMAN, P. C., VAN DIJK, H. K., (1991). A Bayesian routes to unit roots. *Journal of Applied Econometrics*, 6, pp. 387–401.
- SUGITA, K., (2006). Bayesian Analysis of Dynamic Multivariate Models with Multiple Structural Breaks, No 2006-14, Discussion Papers, Graduate School of Economics, Hitotsubashi University, <http://EconPapers.repec.org/RePEc:hit:econdp:2006-14>.
- WANG, J., ZIVOT, E., (2000). A Bayesian time series model of multiple structural changes in level, trend, and variance. *Journal of Business & Economic Statistics*, 18, pp. 374–386.
- ZEILEIS, F., LEISCH, K., H., KLEIBER, C., (2002). Strucchange: An R package for testing for structural change in linear regression models. *Journal of Statistical Software*, 7, pp. 1–38.

APPENDIX

In this appendix, we have derived the posterior probability with the help of likelihood function and prior distribution, which are given below:

(A.1) For alternative hypothesis $H_1 : \rho \in S, \mu_{i1} \neq \mu_{i2}, \sigma_1^2 \neq \sigma_2^2$, expression of posterior probability can be derived with the help of likelihood function (3) and prior distribution given (4), which is given as

$$\begin{aligned}
 & P(y | H_1) \\
 &= \int_l^1 \int_{R_{B+1}^+} \int_{R_{n(B+1)}} L(\rho, \mu, \sigma | y) p(\mu) p(\sigma) p(\rho) d\mu d\sigma d\rho \\
 &= \int_l^1 \int_{R_{B+1}^+} \int_{R_{n(B+1)}} (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \left(\sigma_j^{-n(T_j - T_{j-1})} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1} - (1-\rho)\mu_{ij})^2 \right\} \right] \\
 &\quad \frac{d_1^{c_1}}{\Gamma c_1} \frac{d_2^{c_2}}{\Gamma c_2} \dots \frac{d_{B+1}^{c_{B+1}}}{\Gamma c_{B+1}} (\sigma_1^2)^{-c_1-1} (\sigma_2^2)^{-c_2-1} \dots (\sigma_{B+1}^2)^{-c_{B+1}-1} \exp \left[-\frac{d_1}{\sigma_1^2} \dots - \frac{d_{B+1}}{\sigma_{B+1}^2} \right] (2\pi)^{-\frac{n(B+1)}{2}} \\
 &\quad \sigma_1^{-n} \dots \sigma_{B+1}^{-n} \exp \left[-\frac{1}{2\sigma_1^2} \sum_{i=1}^n (\mu_{i1} - \gamma_{i1})^2 \dots - \frac{1}{2\sigma_{B+1}^2} \sum_{i=1}^n (\mu_{i,B+1} - \gamma_{i,B+1})^2 \right] d\mu d\sigma d\rho \\
 &= \int_l^1 \int_{R_{B+1}^+} \int_{R_{n(B+1)}} \frac{(2\pi)^{\frac{n(T+B+1)}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1})^2 \right. \right. \right. \\
 &\quad \left. \left. - 2(1-\rho) \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1}) \mu_{ij} + (1-\rho)^2 (T_j - T_{j-1}) \sum_{i=1}^n \mu_{ij}^2 + \sum_{i=1}^n \mu_{ij}^2 - 2 \sum_{i=1}^n \gamma_{ij} \mu_{ij} \right. \right. \\
 &\quad \left. \left. \left. + \sum_{i=1}^n \gamma_{ij}^2 + 2d_j \right\} \right\} \right] d\mu d\sigma d\rho \\
 &= \int_l^1 \int_{R_{B+1}^+} \int_{R_{n(B+1)}} \frac{(2\pi)^{\frac{n(T+B+1)}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1})^2 \right. \right. \right. \\
 &\quad \left. \left. + 2d_j + \sum_{i=1}^n \gamma_{ij}^2 + ((1-\rho)^2 (T_j - T_{j-1}) + 1) \sum_{i=1}^n \mu_{ij}^2 \right. \right. \\
 &\quad \left. \left. \left. - 2 \sum_{i=1}^n \left((1-\rho) \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1}) + \gamma_{ij} \right) \mu_{ij} \right\} \right\} \right] d\mu d\sigma d\rho
 \end{aligned}$$

Let us consider

$$A_j = (1 - \rho)^2 (T_j - T_{j-1}) + 1$$

$$B_{ij} = (1 - \rho) \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1}) + \gamma_{ij}$$

Then we may write

$$\begin{aligned} & P(y | H_1) \\ &= \int_l^1 \int_{R_{B+1}^+} \int_{R_{n(B+1)}} \frac{(2\pi)^{\frac{n(T+B+1)}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1})^2 \right. \right. \right. \\ & \quad \left. \left. \left. + \sum_{i=1}^n \gamma_{ij}^2 + 2d_j + A_j \sum_{i=1}^n \left(\mu_{ij} - \frac{B_{ij}}{A_j} \right)^2 - \sum_{i=1}^n \frac{B_{ij}^2}{A_j} \right\} \right] d\mu d\sigma d\rho \\ &= \int_l^1 \int_{R_{B+1}^+} \frac{(2\pi)^{\frac{nT}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]}}{(A_j)^{\frac{n}{2}} \Gamma c_j} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1})^2 \right. \right. \right. \\ & \quad \left. \left. \left. + \sum_{i=1}^n \gamma_{ij}^2 + 2d_j - \sum_{i=1}^n \frac{B_{ij}^2}{A_j} \right\} \right] d\sigma d\rho \\ &= \int_l^1 \int_{R_{B+1}^+} \frac{(2\pi)^{\frac{nT}{2}}}{1-l} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j} (\sigma_j^2)^{\left[\frac{n(T_j - T_{j-1} + 1)}{2} + c_j + 1 \right]}}{(A_j)^{\frac{n}{2}} \Gamma c_j} \right) \exp \left[-\sum_{j=1}^{B+1} \frac{C_{ij}}{\sigma_j^2} \right] d\sigma d\rho \\ &= \int_l^1 \frac{(2\pi)^{\frac{nT}{2}}}{1-l} \prod_{j=1}^{B+1} \frac{d_j^{c_j} \Gamma \left(\frac{n(T_j - T_{j-1} + 1)}{2} + c_j \right)}{\Gamma c_j (A_j)^{\frac{n}{2}} C_j^{\frac{n(T_j - T_{j-1} + 1)}{2} + c_j}} d\rho \end{aligned}$$

where

$$C_{ij} = d_j + \frac{1}{2} \left(\sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} (y_{it} - \rho y_{i,t-1})^2 + \sum_{i=1}^n \gamma_{ij}^2 - \sum_{i=1}^n \frac{B_{ij}^2}{A_j} \right)$$

(A.2) Under unit root hypothesis $H_0 : \rho = 1, \mu_{i1} \neq \mu_{i2}, \sigma_1^2 \neq \sigma_2^2$, the joint likelihood function is given as

$$L(\sigma | y) = \prod_{j=1}^{B+1} L(\sigma_j | y) = \prod_{j=1}^{B+1} \left[(2\pi)^{-\frac{n(T_j - T_{j-1})}{2}} \sigma_j^{-n(T_j - T_{j-1})} \exp \left[-\frac{1}{2\sigma_j^2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} \varepsilon_{it}^2 \right] \right]$$

$$= (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \left(\sigma_j^{-n(T_j - T_{j-1})} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} \Delta y_{it}^2 \right\} \right]$$

By using similar mathematical manipulations as above, we get the posterior probability

$$P(y | H_0)$$

$$= \int_{R_{B+1}^+} L(\sigma | y) p(\sigma) d\sigma$$

$$= \int_{R_{B+1}^+} (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \left(\sigma_j^{-n(T_j - T_{j-1})} \right) \exp \left[-\frac{1}{2} \sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} \Delta y_{it}^2 \right\} \right] \frac{d_1^{c_1}}{\Gamma c_1} \frac{d_2^{c_2}}{\Gamma c_2} \dots \frac{d_{B+1}^{c_{B+1}}}{\Gamma c_{B+1}}$$

$$\left(\sigma_1^2 \right)^{-c_1 - 1} \left(\sigma_2^2 \right)^{-c_2 - 1} \dots \left(\sigma_{B+1}^2 \right)^{-c_{B+1} - 1} \exp \left[-\frac{d_1}{\sigma_1^2} - \frac{d_2}{\sigma_2^2} \dots - \frac{d_{B+1}}{\sigma_{B+1}^2} \right] d\sigma$$

$$= \int_{R_{B+1}^+} (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \left(\frac{d_j^{c_j}}{\Gamma c_j} \left(\sigma_j^2 \right)^{\left[\frac{n(T_j - T_{j-1})}{2} + c_j + 1 \right]} \right) \exp \left[-\sum_{j=1}^{B+1} \left\{ \frac{1}{\sigma_j^2} \left(\frac{1}{2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} \Delta y_{it}^2 + d_j \right) \right\} \right] d\sigma$$

$$= (2\pi)^{-\frac{nT}{2}} \prod_{j=1}^{B+1} \frac{d_j^{c_j} \Gamma \left(\frac{n(T_j - T_{j-1})}{2} + c_j \right)}{\Gamma c_j \left[\frac{1}{2} \sum_{i=1}^n \sum_{t=T_{j-1}}^{T_j} \Delta y_{it}^2 + d_j \right]^{\frac{n(T_j - T_{j-1})}{2} + c_j}}$$